

The temperature and distribution of gas in CL 0016+16 measured with *XMM-Newton*

D.M. Worrall and M. Birkinshaw

Department of Physics, University of Bristol, Tyndall Avenue, Bristol BS8 1TL

2 February 2008

ABSTRACT

We present results of a 37 ks observation of CL 0016+16 with the *XMM-Newton* EPIC instrument. Within 1.5 arcmin of the cluster centre we measure a gas temperature of $kT = 9.13^{+0.24}_{-0.22}$ keV and an abundance of $0.22^{+0.04}_{-0.03}$ times the solar value (1σ uncertainties). This significant improvement over previous measurements has allowed us to revise the estimate of the Hubble constant based on CL 0016+16 to $68 \pm 8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (random error only), close to the value from the Hubble Space Telescope distance-scale project. The total gravitating mass within a radius of 248 kpc of the cluster centre is in good agreement with that found from gravitational lensing over the same region, supporting the assumption of isothermal gas in hydrostatic equilibrium. The gas mass fraction of 0.13 ± 0.02 is in remarkable agreement with that given by cosmological parameters for the Universe as a whole, suggesting that CL 0016+16 is a fair sample of the matter content of the Universe. While there is no spectral or spatial evidence to suggest a cooling flow in CL 0016+16, we find an asymmetrical central X-ray structure which may have a harder spectrum than the cluster as a whole, and be evidence for some merger activity, in addition to a previously reported asymmetry to the west of the cluster. The nearest companion cluster to CL 0016+16 is measured to have a gas temperature of $kT = 3.8^{+0.3}_{-0.3}$ keV and an abundance of $0.6^{+0.3}_{-0.2}$ times the solar value (1σ uncertainties). We also present spectral data for the companion quasar whose spectrum was confused with CL 0016+16 in previous ASCA data.

Key words: cosmology: observations – galaxies: clusters: individual (CL 0016+16, RX J0018.3+1618) – intergalactic medium – quasars: individual: E 0015+162 – X-rays: galaxies: clusters

1 INTRODUCTION

The galaxy content of cluster CL 0016+16 was described by Koo (1981) following its discovery by Kron (1980) in a photometric survey for faint galaxies. Its high redshift, $z = 0.5455$ (Dressler & Gunn 1992), evident richness, high X-ray luminosity ($L_{X, 0.5-4.5 \text{ keV}} \approx 2 \times 10^{45} \text{ ergs s}^{-1}$; White, Silk & Henry 1981), and large Sunyaev-Zel'dovich effect (Birkinshaw, Gull & Moffet 1981) led to extensive work on its properties as the prototypical example of a distant rich cluster.

More recent X-ray and optical observations of the field of CL 0016+16 have found other nearby clusters of galaxies of lower mass but similar redshift (Hughes, Birkinshaw & Huchra 1995; Connolly et al. 1996; Hughes & Birkinshaw 1998a), suggesting that CL 0016+16 is the dominant member of a supercluster at $z \sim 0.5$. A quasar a few arcmin north of CL 0016+16 (Margon, Downes & Spinrad 1983) is also associated with this structure.

The gas properties of CL 0016+16 have been the subject of particular study. Two reasons for this stand out: the intracluster gas can be used to study the cluster mass and baryonic mass fraction, and the combination of X-ray and Sunyaev-Zel'dovich data can be

used to estimate the cluster's distance and hence the Hubble constant or other cosmological parameters.

Grego et al. (2001) used X-ray data and the Sunyaev-Zel'dovich effect of CL 0016+16 to measure a gas mass fraction, f_g , that is close to the value seen in nearby clusters, implying that there is little evolution in f_g between $z \sim 0.5$ and today. If cluster gas is the dominant baryonic component of CL 0016+16, and CL 0016+16 is a fair sample of the matter content of the Universe, f_g will be close to the baryonic mass fraction of the Universe, Ω_b/Ω_m . However, the temperature of the cluster gas is a necessary component of the calculation of both the gas and total masses, and improved measurements of the gas temperature would be expected to improve the reliability of cosmological conclusions drawn from studies of the mass components of the cluster.

Good knowledge of the cluster temperature would also allow a comparison between the mass derived from the assumption that the cluster gas is in hydrostatic equilibrium and the mass measured by Smail et al. (1997) using a gravitational shear technique. This could set limits on the three-dimensional shape of CL 0016+16, which is a source of systematic error in the use of any cluster as a cosmological tracer.

Three independent Sunyaev-Zel'dovich effect measurements of CL 0016+16 (Hughes & Birkinshaw 1998b; Reese et al. 2000; Grainge et al. 2002) have been combined with X-ray data to measure the distance of the cluster. All three lead to values for the Hubble constant of about $47 \text{ km s}^{-1} \text{ Mpc}^{-1}$ assuming $q_0 = \frac{1}{2}$ (this would be $57 \text{ km s}^{-1} \text{ Mpc}^{-1}$ for a cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$). The low value of the Hubble constant found by this technique might be attributed to projection or selection effects, calibration uncertainties, unresolved substructure, or some other problem, but it is clear that a substantial fraction of the *random* component of the error is due to uncertainty in the X-ray temperature. A deep X-ray observation with *XMM-Newton*, which can detect arriving cluster photons of energy up to about 10 keV, and so can make a better measurement of the temperature of a hot cluster such as CL 0016+16 (which Hughes & Birkinshaw 1998b found to have $kT \approx 7.55 \text{ keV}$ in the rest frame), would be expected to resolve this problem.

Thus we obtained a long *XMM-Newton* observation of CL 0016+16 with the aim of improving the measurement of the temperature and metal abundance of the cluster, and hence making it more useful as a cosmological tracer. This paper describes the data, their analysis, and the implications for the temperature and structure of CL 0016+16 and one of the companion clusters in the supercluster. We also present the *XMM-Newton* spectrum of the companion quasar. We revise the estimate of the Hubble constant based on CL 0016+16 using our new temperature for the cluster, and compare an estimate of the total gravitating mass with that measured from gravitational lensing. We compare the gas mass fraction in CL 0016+16 with the value for the Universe as a whole given by Ω_b/Ω_m . Finally, we point out structural features of CL 0016+16 which require further study.

We use a cosmology in which $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$, and we adopt $H_o = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ outside the discussion of the use of CL 0016+16 as a tracer of the Hubble flow.

2 OBSERVATIONS

XMM-Newton observed CL 0016+16 on December 30th, 2000. In this paper we concentrate on data from the European Photon Imaging Camera (EPIC) which incorporates one pn (Sröder et al. 2001) and two MOS (MOS1 and MOS2; Turner et al. 2001) CCD-array cameras. The cameras were operated in Full Frame mode with the medium optical blocking filter. The data were provided to us in the form of two separate observations, of maximum livetime (for the MOS cameras) roughly 31.3 ks and 5.5 ks, separated by 110 minutes. Analysis presented here uses the latest software available from the *XMM-Newton* project at <http://xmm.vilspa.esa.es/>, SAS v5.3.3. Calibrated event files were generated using the EMCHAIN and EPCHAIN scripts (described in detail on the project WWW site) and, after merging the data for each camera separately for the two observations, we have selected for analysis the good events with patterns 0 to 12 from the MOS data and patterns 0 to 4 from the pn data.

The light curves extracted from the full-field, pattern=0, 10–12 keV events were examined for intervals of high particle background, which we defined as those with $> 0.1 \text{ ct s}^{-1}$ for MOS1 and MOS2, and $> 0.2 \text{ ct s}^{-1}$ for the pn. After removing the high-background intervals, the remaining exposure times were roughly 35.9 ks, 36 ks, and 24.3 ks for MOS1, MOS2, and pn, respectively.

Our analysis follows the prescription in Arnaud et al. (2002). This is superior to the simple use of local background in analysing




Figure 1. Combined MOS1, MOS2 and pn data for the observation of CL 0016+16 in the 0.3–5 keV energy band, with a logarithmic intensity scale, a pixel size of 3 arcsec, and showing J2000 equatorial coordinates. Chip boundaries are visible. This image is vignetting corrected and without background subtraction. The bright source at a declination of roughly $16^\circ 29' 26''$, north of CL 0016+16, is the quasar E 0015+162 (Margon et al. 1983), and the poor companion cluster RX J0018.3+1618 (Hughes et al. 1995) is visible to the SW.

extended-source data because it treats separately the non-vignetted particle background and the vignetted cosmic X-ray background. First the EVIGWEIGHT task was run on both CL 0016+16 and blank-sky data so that after their subtraction the spectra and surface-brightness profiles (with local background) that are extracted are corrected for vignetting. The vignetting-corrected MOS1, MOS2 and pn event files have been combined for display purposes. Figure 1 shows the 0.3–5 keV image of a wide field, centred on CL 0016+16, from this merged event file.

We use the data compiled by Lumb (2002), taken from deep-sky exposures cleaned of sources, to subtract from our data the particle background, which is assumed to provide all the pattern=0 counts between 10 and 12 keV. After applying our good-time-interval screening, the full-field, pattern=0, 10–12 keV count rates are the same in the CL 0016+16 and blank-sky fields to within 1%, 7% and 16%, for the MOS1, MOS2, and pn, respectively. The relevant scaling factor is applied to the vignetting-corrected blank-sky data before subtraction from the image data.

The blank-sky data are not expected to be a good representation of the background in the CL 0016+16 data at low energies. Here the cosmic X-ray background is the dominant contributor and, more importantly, at energies below about 1 keV the medium optical blocking filter used for our observations has a lower transparency than the thin filter used for the blank-sky data. This results in an over-subtraction of background at the lowest energies. This is corrected for by using a blank-sky-subtracted source-free region of the CL 0016+16 image as a background for the blank-sky-subtracted on-source data. By selecting this background region to be an annulus of radii 4 and 11 arcmin, between position angles 90 and 120 degrees, and 220 and 310 degrees, we limited the number of source-contaminated exclusion regions to one circle of radius 30 arcsec.

As data have been corrected for vignetting, spectral fitting uses on-axis response matrices and effective areas. We have used the version 6 response matrices made available by the project, and have

generated on-axis effective areas using the ARFGEN task. Bins between energies of 0.4 and 10 keV are included in spectral fitting using the XSPEC software.

3 EMISSION-WEIGHTED CLUSTER TEMPERATURE

For good signal to noise, we have found the cluster spectral parameters using the net counts (roughly 32,600 counts between 0.4 and 10 keV) extracted from a circle of radius 1.5 arcmin. We fitted the data to a redshifted MEKAL model modified by local Galactic absorption. We let the relative normalization between the MOS1, MOS2 and pn data be free but all other parameters are in common. This results in MOS1 and MOS2 normalizations in good agreement, but the pn normalization is a few per cent lower, consistent with the evaluation of the EPIC flux calibration by Saxton (2002).

We obtain a value of $N_H = (4.3 \pm 0.6) \times 10^{20} \text{ cm}^{-2}$ (95% confidence for one interesting parameter) for the Galactic hydrogen column density when it is a free parameter, in agreement with the value of $4.04 \times 10^{20} \text{ cm}^{-2}$ found from 21 cm observations (Dickey & Lockman 1990). The value of the redshift fitted from the X-ray spectrum, $z = 0.532^{+0.011}_{-0.013}$, is slightly low as compared with published optical values of 0.5455 (Dressler & Gunn 1992) and 0.5481 (Ellingson et al. 1998), suggesting that, although the EPIC energy scale is believed to be calibrated to better than 5 eV (Kirsch 2002), there may be a small additional systematic error in the CL 0016+16 observations. We have fixed the Galactic absorption to the 21 cm value but have allowed the cluster redshift to be free in finding the cluster parameters and uncertainties reported in this paper.

The best fit gives $kT = 9.13^{+0.24}_{-0.22} (+0.49_{-0.44})$ keV and an abundance of $0.22^{+0.04}_{-0.03} (+0.07_{-0.065})$ times the solar value, where uncertainties are statistical and quoted as 1σ (unbracketed), with 95% errors given in brackets, for one interesting parameter. The model is shown together with the data in Figure 2, and joint-confidence uncertainty contours are shown in Figure 3. With a reduced χ^2 of 1.09 ($\chi^2 = 859$ for 786 degrees of freedom) the null hypothesis probability is only 4%, but this is likely to result from the high statistical precision of the data coupled with small remaining systematic calibration uncertainties (Saxton 2002). An apparently random scatter in the residuals to the pn data is diminished if the pattern=0 events only are used for the pn, making the combined MOS and pn fit acceptable (reduced χ^2 of 1.01: $\chi^2 = 706$ for 697 degrees of freedom). However, as the best-fit parameter values are unchanged, we elect to include patterns 0–4 in the fits to improve statistical accuracy.

There is consistency between the parameter values fitted separately for data from the MOS ($kT = 9.3^{+1.1}_{-0.7}$, abundance = $0.22^{+0.09}_{-0.08}$ times the solar value, $z = 0.534^{+0.015}_{-0.013}$, $N_H = 4.4^{+0.9}_{-1.0} \times 10^{20} \text{ cm}^{-2}$) and pn ($kT = 8.6^{+0.8}_{-0.6}$, abundance = $0.22^{+0.1}_{-0.09}$ times the solar value, $z = 0.525^{+0.02}_{-0.03}$, $N_H = 4.3^{+0.7}_{-0.8} \times 10^{20} \text{ cm}^{-2}$), all quoted with 95% uncertainties for one interesting parameter.

The observed 2–10 keV flux within the circle of 1.5 arcmin radius is measured to be $1.43 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and $1.39 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$ with the MOS and pn cameras, respectively. Between 0.4 and 10 keV the values are $2.24 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and $2.18 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$, respectively. The average volume-weighted emission measures ($\int n_e n_p dV / 4\pi D_L^2$) are $(1.22 \pm 0.02) \times 10^{11} \text{ cm}^{-5}$ and $(1.18 \pm 0.02) \times 10^{11} \text{ cm}^{-5}$, respectively (1σ uncertainties)¹.




Figure 2. XMM-Newton EPIC spectrum of CL 0016+16 from counts within a circle of radius 1.5 arcmin. The upper spectrum is from the pn, and the lower spectrum is an overlay of MOS1 and MOS2 data points. The fit is to an isothermal model with $kT = 9.1$ keV and an abundance of 0.22 times the solar value.




Figure 3. CL 0016+16 spectral-parameter uncertainty contours using counts within a circle of radius 1.5 arcmin. Contours are 1σ , 90% and 99% for two interesting parameters, with the normalizations and redshift allowed to vary.

4 GAS DISTRIBUTION

The intensity distribution of X-ray emission in the central part of CL 0016+16 (Fig. 4) shows an apparent filamentary structure in the shape of an inverted ‘V’. A true colour image, made using software from the Chandra project (the CIAO DMIMG2.JPG task), with counts in 0.3–1 keV, 1–2 keV, and 2–8 keV corresponding to red, green and blue, respectively, suggests that the emission at the centre of the ‘V’ may be hotter than in the surrounding part of the cluster. However the prong to the SE appears cooler than average.

These spectral differences also appear when we use the IRAF/STSDAS ELLIPSE program to model the underlying large-

thors, the normalization given by fitting thermal plasma models is $10^{-14} \times (1+z)^2$ times the volume-weighted emission measure.

¹ In the XSPEC X-ray spectral-fitting software used here, and by most au-




Figure 4. Same as Figure 1, showing only the centre of CL 0016+16. Scale is linear with black corresponding to about 70 counts and the faintest pixels containing about 20 counts. An inverted V-shaped filamentary structure is apparent. This is also seen in the individual MOS and pn images, and is unrelated to chip orientation.

scale structure of the cluster, and then subtract a scaled version of the model first from a 0.3–1.5 keV image and then a 2–8 keV image. The most prominent positive residuals in the soft image track the SE prong of the inverted ‘V’, whereas in the hard image they are at the cluster centre. The diffuse asymmetric emission to the west of the cluster, at about 1.5 arcmin from the centre (Fig. 1), appears somewhat softer than CL 0016+16 as a whole. This western diffuse emission was previously discussed by Neumann & Böhringer (1997) using *ROSAT* data, and is interpreted as a merging subcomponent of the cluster that is associated with a peak in the weak-lensing mass map of Smail et al. (1995).

We have fitted separately the counts within a source-centred circle of radius 0.6 arcmin and a source-centred annulus of radii 0.6 and 1.5 arcmin using the method and model described in §3. The results (Fig. 5) indeed show that the inner region is hotter and of higher abundance than the outer one, at a combined confidence of 99% for two interesting parameters. The evidence for spatial and temperature structure in the inner regions may result from merger activity. A deeper observation would be required to probe this in detail.

We have extracted the circularly-symmetric radial profile of background-subtracted 0.3–5 keV counts and fitted it to a spherical, isothermal, β model of surface brightness $b_x(\theta) \propto (1 + [\theta/\theta_c]^2)^{-3\beta+0.5}$, convolved with the point spread function (PSF). While the cluster is clearly ellipsoidal in appearance, and displays hotter regions within its core, the value of β and the average core radius of the model produced by fits of spherical functions are close to those obtained by making more complicated ellipsoidal fits (see Tables 1 and 2 in Hughes & Birkinshaw 1998b). If the fit results are to be used in the calculation of the distance of the cluster (as, for example, in §7), then values obtained from ellipsoidal fits could not be used directly to estimate H_0 , since the projection is unknown for any individual cluster. However, H_0 estimates obtained from spherical model fits for a *population* of clusters are almost unbiased, provided that the clusters are selected in an orientation-unbiased fashion. Thus spherical modeling of clusters is sufficient for the purpose of building up a sample of distance measures from clusters.

To construct the radial profile we have summed the counts from the three cameras. For the PSF we have used the on-axis pa-




Figure 5. Same as Figure 3, separating counts within a circle of radius 0.6 arcmin (upper right contours) from those within an annulus of radii 0.6 and 1.5 arcmin.

rameter values for 1 keV from Ghizzardi’s (2001, and private communication in 2002) fits of a β model to data from stellar sources. We have summed the MOS1, MOS2, and pn PSFs, weighted by each camera’s contribution to the counts from CL 0016+16. The counts from a circle of radius 20 arcsec around the quasar 3.25 arcmin to the north of CL 0016+16 (Fig. 1) are excluded from the fits.

The radial profile and best-fit model are shown in Figure 6. While the filamentary structure seen in Figure 4 leads to a poor fit within about 20 arcsec of the centre, this has negligible effect on the values found for β and θ_c , and the overall fit is acceptable ($\chi^2 = 50.3$ for 37 degrees of freedom). The values found are $\beta = 0.697 \pm 0.010$, and $\theta_c = 36.6 \pm 1.1$ arcsec, where, since the parameters are strongly correlated, the errors quoted are one sigma for two interesting parameters. Nothing in the radial profile or spectral properties of CL 0016+16 is suggestive of a cooling flow.

5 COMPANION CLUSTER, RX J0018.3+1618

The companion cluster to the SW of CL 0016+16 (Fig. 1) appears redder than CL 0016+16 in a true colour image (made as in §4). We have found spectral-parameter values using on-source net counts from a circle of radius 1.5 arcmin (roughly 3,200 counts between 0.4 and 10 keV) in a similar manner to the extraction and fitting for CL 0016+16 itself. The fit to a MEKAL model is good, with $\chi^2 = 109$ for 101 degrees of freedom. The column density of $N_H = 5.8^{+2.6}_{-2.3} \times 10^{20} \text{ cm}^{-2}$ (1σ errors) is consistent with the value from 21 cm measurements ($4.09 \times 10^{20} \text{ cm}^{-2}$; Dickey & Lockman 1990), and so we fix the value to the radio-derived measurement. Figure 7 shows the data and model whose best-fit parameter values and uncertainties are as follows, where the latter are 1σ (95%) confidence for one interesting parameter where all the other parameter values are allowed to vary: $kT = 3.8^{+0.3}_{-0.3} (^{+0.8}_{-0.6})$, abundance = $0.6^{+0.3}_{-0.2} (^{+0.6}_{-0.4})$ times solar, $z = 0.472^{+0.062}_{-0.022} (^{+0.09}_{-0.05})$. The redshift is consistent (at 95% confidence, or 1σ if the systematic error of 0.016 suggested for CL 0016+16 is applied) with the value of 0.5506 ± 0.0012 found from optical measurements by Hughes et al. (1995) who were able only to place a lower limit on the cluster temperature ($kT > 1.5$ keV in the observed frame) using data from *ROSAT*. The precision with which the spectral parameters for the




Figure 6. 0.3–5 keV radial profile of CL 0016+16 after subtraction of deep-sky and local backgrounds, as described in §2. The curve is the best fit circularly symmetric β model convolved with the PSF: $\beta = 0.697$, $\theta_c = 36.546$ arcsec. The horizontal dotted line shows the contributions, taken into account in the fitting, of the model to the local background annulus.




Figure 7. XMM-Newton EPIC spectrum of the companion cluster from counts within a circle of radius 1.5 arcmin. The upper spectrum is from the pn, and the lower spectrum is an overlay of MOS1 and MOS2 data points. The fit is to an isothermal model with $kT = 3.8$ keV and an abundance of 0.6 times the solar value.

companion cluster are measured is similar to that with which the parameters for CL 0016+16 were known prior to this work (Furuzawa et al. 1998; Hughes & Birkinshaw 1998b).

The net observed 2–10 keV and 0.4–10 keV fluxes are measured to be 7.7×10^{-14} ergs cm $^{-2}$ s $^{-1}$ and 1.6×10^{-13} ergs cm $^{-2}$ s $^{-1}$, respectively. The volume-weighted emission measure is $(1.09 \pm 0.12) \times 10^{10}$ cm $^{-5}$.

Because this cluster lies on a gap between CCDs in the pn




Figure 8. XMM-Newton EPIC spectrum of the companion quasar from counts within a circle of radius 40 arcsec. The upper spectrum is from the pn, and the lower spectrum is an overlay of MOS1 and MOS2 data points. The fit is to a power law of photon spectral index 2.38, with a narrow Fe fluorescence line of rest energy 6.4 keV.

camera, as can be seen in Figure 1, we have fitted the radial profile using only data from the MOS cameras. The parameter values found for a β model, convolved with the PSF corresponding to the off-axis position of the cluster, are of comparable value and precision to the values found by Hughes et al. (1995) using ROSAT PSPC data. We find $\beta = 0.74^{+0.36}_{-0.17}$, $\theta_c = 28^{+15}_{-9}$ arcsec (1σ uncertainties).

6 COMPANION QUASAR, E 0015+162

The companion radio-quiet quasar of redshift 0.554 (Margon et al. 1983) to the north of CL 0016+16 (Fig. 1) has a relatively steep spectrum, and we use the net counts between 0.3 and 10 keV within a source-centred circle of radius 40 arcsec for our fits. A fit to an absorbed power law is acceptable ($\chi^2 = 113$ for 126 degrees of freedom), with the column density consistent with the Galactic value from 21 cm measurements. χ^2 is reduced by 7.1 for one extra parameter (significant at 99% confidence on an F test) when a narrow Fe fluorescence line, modeled as a Gaussian of fixed rest-frame energy 6.4 keV and $\sigma = 10$ eV (small compared with the roughly 130 eV FWHM resolution of EPIC at the observed energy) is included in the model. If the line energy is allowed to be a free parameter, the best-fit value and 1σ error is $6.1^{+0.6}_{-0.1}$. Fixing the line energy at 6.4 keV, the following parameter values and uncertainties are found, where the latter are 1σ (95%) confidence for one interesting parameter where all the other parameter values are allowed to vary: power-law photon index $\Gamma = 2.38 \pm 0.06(\pm 0.13)$, $N_H = 3.4 \pm 1.0(\pm 1.9) \times 10^{20}$ cm $^{-2}$, equivalent width of the Fe line = $308^{+203}_{-123}({}^{+352}_{-271})$ eV. The spectrum is shown in Figure 8. The observed 0.5–10 keV flux from the quasar is $1.61 \pm 0.07 \times 10^{-13}$ ergs cm $^{-2}$ s $^{-1}$ (1σ error) and the luminosity is 2.15×10^{44} ergs s $^{-1}$.

The AGN is roughly 8% the strength of CL 0016+16 (in a cluster-centred circle of radius 4 arcmin) in terms of net counts between 0.4 and 10 keV (roughly 3,500 counts from the AGN). Reeves et al. (1997) fitted the ASCA spectrum of the AGN and found only an upper limit for narrow Fe fluorescence-line emission,

but the ASCA spectrum would have been heavily contaminated with emission from CL 0016+16, and this would seem to account for the inconsistency between our results and the flat spectral index (1.99 ± 0.09) and large flux (2.92×10^{-12} ergs cm $^{-2}$ s $^{-1}$ between 0.5 and 10 keV) found by these authors. Our spectral index is consistent with the value of 2.54 ± 0.18 (1σ error) found for the quasar by Hughes & Birkinshaw (1998b) from a combined thermal and power-law model fitted jointly to ASCA and ROSAT PSPC data in order to characterize the emission both from CL 0016+16 and the quasar. Their power-law normalization is also consistent with our better-determined value. *XMM-Newton* has improved the precision with which the spectral index is measured by a factor of three, and has provided the first evidence for an Fe fluorescence line in this radio-quiet quasar.

7 HUBBLE CONSTANT AND GAS CONTENT OF CL 0016+16

XMM-Newton has measured the temperature of CL 0016+16's atmosphere to $\pm 2.5\%$. This is a considerable improvement on the 9% uncertainty obtained using ASCA and ROSAT (Hughes & Birkinshaw 1998b), and so reduces one of the main sources of random error in the Hubble constant as deduced by comparing the X-ray and Sunyaev-Zel'dovich effect properties of the cluster.

The *XMM-Newton*-derived emission measure within 1.5 arcmin of the centre, $(1.20 \pm 0.03) \times 10^{11}$ cm $^{-5}$, from the weighted average of the MOS and pn values, implies an integrated cluster emission measure of

$$\frac{\int n_e n_p dV}{4\pi D_L^2} = (1.75 \pm 0.05) \times 10^{11} \text{ cm}^{-5} \quad (1)$$

if the fraction of the emission contained within a 1.5-arcmin radius cylinder is extrapolated to the full cluster using the best-fit radial profile parameters. This emission measure is related to the angular diameter distance of the cluster, D_A , and the central electron density, n_{e0} , by

$$\text{EM} = \frac{D_A \theta_c^3 n_{e0}^2}{\eta (1+z)^4} \frac{\sqrt{\pi}}{4} \frac{\Gamma(3\beta - \frac{3}{2})}{\Gamma(3\beta)} \quad (2)$$

if the *XMM-Newton*-derived β model is used to describe the overall structure of the cluster, and so the X-ray data provide a measurement of $D_A n_{e0}^2$. The symbol η in equation (2) is the electron/proton ratio, ≈ 1.17 . The emission measure (1) is less than half the value given by Hughes & Birkinshaw (1998b), because of an error in the way that plasma thermal emission normalizations were reported by XSPEC. Since the emission measure was not used in the calculation of the cluster distance in the Hughes & Birkinshaw paper, however, the conclusions about the cluster distance there are unchanged by this error.

The β model can also be used to calculate the central Sunyaev-Zel'dovich effect for the cluster,

$$\Delta T_0 = -2 T_{\text{rad}} \sigma_T \left(\frac{kT}{m_e c^2} \right) n_{e0} D_A \theta_c \sqrt{\pi} \frac{\Gamma(\frac{3}{2}\beta - \frac{1}{2})}{\Gamma(\frac{3}{2}\beta)} \quad (3)$$

where T_{rad} is the thermodynamic temperature of the microwave background radiation and σ_T is the Thompson scattering cross-section.

The central Sunyaev-Zel'dovich effect of CL 0016+16 is found by fitting a one-dimensional profile (Hughes & Birkinshaw 1998b) or a two-dimensional map (Reese et al. 2000; Grainge et

al. 2002) of the effect. Using slightly different β models for the cluster atmosphere, values $\Delta T_0 = -1.20 \pm 0.19$, -1.24 ± 0.11 , and -1.08 ± 0.11 mK were found in these investigations. We re-analysed the Hughes & Birkinshaw (1998b) data using the β -model parameters based on our *XMM-Newton* data, and scaled the interferometric data (which are relatively insensitive to the β -model parameters along degeneracy lines in the $[\beta, \theta_c]$ plane) to these same parameters. A weighted combination of all three independent values leads to a central Sunyaev-Zel'dovich effect for CL 0016+16 of $\Delta T_0 = -1.26 \pm 0.07$ mK. Via equation (3), this implies a measurement of $D_A n_{e0}$.

Combining our measurements of $D_A n_{e0}^2$ and $D_A n_{e0}$, and using the optical cluster redshift of 0.5481 (Ellingson et al. 1998), we find the angular diameter distance and central electron density of CL 0016+16 to be

$$D_A = 1.36 \pm 0.15 \text{ Gpc, and} \quad (4)$$

$$n_{e0} = (8.8 \pm 0.5) \times 10^{-3} \text{ cm}^{-3}. \quad (5)$$

In a Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, this corresponds to a Hubble constant of 68 ± 8 km s $^{-1}$ Mpc $^{-1}$ (random error only), which is about 45% larger than the values estimated by Hughes & Birkinshaw (1998b), Reese et al. (2000), and Grainge et al. (2002). The change in H_0 arises about equally from the increased best-fit gas temperature found in the present *XMM-Newton* study and the change from a flat matter-dominated cosmology to a flat Λ CDM cosmology.

In addition to the statistical error in equation (4), the distance is subject to systematic errors, the largest of which are 5% calibration uncertainties for the Sunyaev-Zel'dovich effect and X-ray normalization, a 20% uncertainty from possible projection effects, and a 15% uncertainty from substructure within the cluster (Hughes & Birkinshaw 1998b). Overall, therefore, we deduce an angular diameter distance for CL 0016+16 of 1.36 ± 0.15 (statistical) ± 0.34 (systematic) Gpc, which implies a Hubble constant of 68 ± 8 (statistical) ± 18 (systematic) km s $^{-1}$ Mpc $^{-1}$. This is consistent with the value of the Hubble constant found in the HST distance-scale project (Mould et al. 2000), but the systematic errors are too large for a useful independent measurement of Ω_Λ to result from our distance for CL 0016+16.

It is interesting to use the X-ray data for CL 0016+16 to estimate the mass of the cluster within the region well-probed by *XMM-Newton*. For an isothermal gas in hydrostatic equilibrium in a spherical cluster, the total gravitating mass within a cylinder of radius r about the cluster centre is

$$M_{\text{tot}}(r) = \frac{3\pi\beta}{2} \frac{kT}{\mu m_p G} \frac{r^2}{r_c} \left(1 + \frac{r^2}{r_c^2} \right)^{-\frac{1}{2}}. \quad (6)$$

For $r = 248$ kpc, this corresponds to a mass of $(2.0 \pm 0.1) \times 10^{14} M_\odot$. This value is consistent with the mass of $(2.7 \pm 0.9) \times 10^{14} M_\odot$ measured from the gravitational lensing effect of the cluster within the same cylindrical radius (Smail et al. 1997; value converted to our adopted cosmology).

We can compare this total gravitating mass with the gas mass within the same cylindrical radius, based on the central electron density that we measure. The result, $M_{\text{gas}}(248 \text{ kpc}) = (2.6 \pm 0.2) \times 10^{13} M_\odot$ indicates that X-ray emitting gas contributes a fraction (0.13 ± 0.02) of the total mass of the cluster in this 248-kpc radius cylinder (for $H_0 = 70$ km s $^{-1}$ Mpc $^{-1}$), or (0.13 ± 0.04) independent of cosmology if we use the cluster distance in equation (4) and the electron density in equation (5).

The measured gas mass fraction is consistent with the value

$0.14^{+0.09}_{-0.04}$ within a radius of 65 arcsec (415 kpc) which is obtained by correcting the result of Grego et al. (2001) to the cosmology that we use and cluster temperature that we measure. Our temperature makes CL 0016+16's gas mass fraction more consistent with the average in the Grego et al. sample, and with the averages for the Myers et al. (1997) and Mohr et al. (1999) samples. Since these samples derived the gas mass fraction using different analyses of the available X-ray and/or Sunyaev-Zel'dovich effect data, their agreement about the gas mass fraction and its lack of change with redshift suggests that the assumptions used, notably the lack of small-scale density or thermal structure in the atmospheres, are not seriously in error. Since the baryonic matter content of the X-ray gas dominates that of galaxies in clusters, and $\Omega_b/\Omega_m = 0.12 \pm 0.02$ (Turner 2002) for the Universe as a whole, it appears that our 248-kpc radius cylinder through CL 0016+16, and clusters of galaxies in general, are fair samples of the matter content of the Universe. Studies of how the gas mass fraction in clusters varies with redshift are therefore unlikely to provide useful information on cluster evolution, at least at the current level of accuracy of the measurements.

8 SUMMARY

With the EPIC camera of *XMM-Newton* we have measured the temperature and distribution of the intra-cluster medium of CL 0016+16 to unprecedented precision. The emission-weighted temperature within 1.5 arcmin of the cluster centre is $kT = 9.13^{+0.24}_{-0.22}$ keV and the abundance is $0.22^{+0.04}_{-0.03}$ (1σ uncertainties). The higher best-fit temperature than found in previous work using ASCA (Furuzawa et al. 1998; Hughes & Birkinshaw 1998b) affects the cluster's position on the temperature-luminosity plane, now moving it into good agreement with the $L_x - T$ relation for nearby clusters (see figure 7 of Schindler 1999).

Although there is some ellipticity in the plane of the sky, as previously reported based on ROSAT data (Neumann & Böhringer 1997; Hughes & Birkinshaw 1998), we find an acceptable fit of the overall envelope out to 5 arcmin (1.9 Mpc) to a circularly-symmetric β model, representative of gas in hydrostatic equilibrium, with $\beta = 0.697 \pm 0.010$, and core radius $\theta_c = 36.6 \pm 1.1$ arcsec (where errors are 1σ for two interesting parameters since β and θ_c are strongly correlated).

While there is no spectral or spatial evidence to suggest a cooling flow in CL 0016+16, an asymmetric enhancement to the west of the cluster is apparent in the *XMM-Newton* image. Such an asymmetry was previously reported by Neumann & Böhringer (1997) and interpreted as a merging subcomponent of the cluster. The *XMM-Newton* data also show a more central asymmetric X-ray structure which may have a harder spectrum than the cluster as a whole, and be evidence of merger activity. A deeper X-ray exposure would be required to probe this in more detail.

The improved spectral and spatial data provided by *XMM-Newton* have allowed us to reduce the random error on the angular diameter distance of CL 0016+16 to 11%. The resulting estimate of the value of the Hubble constant, $68 \pm 8(\text{random}) \pm 18(\text{systematic}) \text{ km s}^{-1} \text{ Mpc}^{-1}$, is in good agreement with the value $71 \pm 6 \text{ km s}^{-1} \text{ Mpc}^{-1}$ from the HST distance-scale project (Mould et al. 2001). The largest remaining systematic component of the error in our estimate of the Hubble constant arises from projection and substructure effects, and could be reduced by a factor ~ 3 by applying the technique to a sample of 10 – 20 clus-

ters with equally-good X-ray and Sunyaev-Zel'dovich effect data as CL 0016+16.

The total gravitating mass within a radius of 248 kpc of the cluster centre is in good agreement with that found from gravitational lensing over the same region (Smail et al. 1997), which supports the assumption that the gas in CL 0016+16 can be treated as isothermal and in hydrostatic equilibrium. The measured gas mass fraction of 0.13 ± 0.02 is in remarkable agreement with that given by the cosmological parameters for the Universe as a whole, suggesting that CL 0016+16 is a fair sample of the matter content of the Universe.

The cluster CL 0016+16 is believed to be the dominant member of a supercluster at $z \sim 0.5$. The nearest companion cluster and relatively bright companion quasar E 0015+162 (which is of an X-ray luminosity more typical of a local Seyfert galaxy) are both well measured in the *XMM-Newton* EPIC data. In previous ASCA data the emission from the quasar is confused with that from CL 0016+16 and so values reported for the quasar based on that observation are unreliable. We find that the spectrum of E 0015+162 fits well a power law absorbed only by the column density of gas through our Galaxy, with a relatively steep power-law photon index of $\Gamma = 2.38 \pm 0.06$. There is evidence for the detection of an Fe fluorescence line.

XMM-Newton finds that the spectrum of the nearest companion cluster, RX J0018.3+1618, fits a thermal model with $kT = 3.8^{+0.3}_{-0.3}$ keV and abundance $= 0.6^{+0.3}_{-0.2}$ times solar. These uncertainties for RX J0018.3+1618 are comparable to those of earlier measurements of CL 0016+16. Deeper and broader X-ray observations of the supercluster may provide important information on a massive structure at an early phase of its evolution.

ACKNOWLEDGEMENTS

This work is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and the USA (NASA). The observation was made as part of the *XMM-Newton* Survey Science Centre (SSC) Guaranteed-Time programme. We thank Mike Watson for organizing the SSC and its scientific programme.

REFERENCES

- Arnaud, M., et al., 2002, A&A, in press (astro-ph/0204306)
- Birkinshaw, M., Gull, S.F., Moffett, A.T., 1981, ApJ, 251, L69
- Connolly, A.J., Szalay, A.S., Koo, D.C., Romer, A.K., Holden, B., Nichol, R.C., Miyaji, T., 1996, ApJ, 473, L67
- Dickey, J.M., Lockman, F.J., 1990, ARA&A, 28, 215
- Dressler, A., Gunn, J.E., 1992, ApJS, 78, 1
- Ellingson, E., Yee, H.K.C., Abraham, R.G., Morris, S.L., Carlberg, R.G., 1998, ApJS, 116, 247
- Furuzawa, A., Tawara, Y., Kunieda, H., Yamashita, K., Sonore, T., Tanaka, Y., Mushotzky, R., 1998, ApJ, 504, 35
- Ghizzardi, S. 2001, 'In flight calibration of the PSF for the MOS1 and MOS2 cameras', EPIC-MCT-TN-001, available from <http://xmm.vilspa.esa.es>
- Grainge, K., Grainger, W.F., Jones, M.E., Kneissl, R., Pooley, G.G., Saunders, R., 2002, MNRAS, 329, 890
- Grego, L., Carlstrom, J.E., Reese, E.D., Holder, G.P., Holzapfel, W.L., Joy, M.K., Mohr, J.J., Patel, S., 2001, ApJ, 552, 2
- Hughes, J.P., Birkinshaw, M., 1998a, ApJ, 497, 645
- Hughes, J.P., Birkinshaw, M., 1998b, ApJ, 501, 1
- Hughes, J.P., Birkinshaw, M., Huchra, J.P., 1995, ApJ, 448, L93

- Kirsch, M.G.F. 2002, presentation to the *XMM-Newton* SAS Workshop on May 7th, 2002, available from <http://xmm.vilspa.esa.es>
- Koo, D.C., 1981, *ApJ*, 251, L75
- Kron, R.G., 1980, PhD thesis, University of California, Berkeley
- Lumb, D. 2002, 'EPIC background files', XMM-SOC-CAL-TN-0016, available from <http://xmm.vilspa.esa.es>
- Margon, B., Downes, R.A., Spinrad, H., 1983, *Nature*, 301, 221
- Mohr, J.J., Mathiesen, B. & Evrard, A.E., 1999, *ApJ*, 517, 627
- Mould, J.R. et al., 2000, *ApJ*, 529, 786
- Myers, S.T. et al., 1997, *ApJ*, 485, 1
- Neumann, D.M., Böhringer, H., 1997, *MNRAS*, 289, 123.
- Reese, E.D., et al., 2000, *ApJ*, 533, 38
- Reeves, J.N., Turner, M.J.L., Ohashi, T., Kii, T., 1997, *MNRAS*, 292, 468
- Saxton, R.D., 2002, 'A statistical evaluation of the EPIC flux calibration', XMM-SOC-CAL-TN-0023, available from <http://xmm.vilspa.esa.es>
- Schindler, S., 1999, *A&A*, 349, 435
- Smail, I., Ellis, R.S., Fitchett, M.J., Edge, A.C., 1995, *MNRAS*, 273, 277
- Smail, I. et al., 1997, *ApJ*, 479, 70
- Stüder, L. et al., 2001, *A&A*, 365, L18
- Turner, M.J.L. et al., 2001, *A&A*, 365, L27
- Turner, M.S., 2002, *ApJ*, 576, L101
- White, S.D.M., Silk, J.I., Henry, J.P., 1981, *ApJ*, 251, L65